

Table 15. IBM RS/6000 590 Timings^(a)Ethylene, 16 electrons, ¹A₁ (D_{2h}), Basis Set=6-311++G**(74 functions, 6-term d's)^(b)

Method	Gaussian 92 (E)	Gaussian 92/DFT(G)	Gaussian 94 (C)
Conv. RHF	1/12 (23)	1/11 (19)	1/14 (30)
Direct RHF	4/36 (42)	3/31 (42)	4/36 (43)
RHF Gradient	9/21 (30)	8/19 (36)	12/26 (45)
RHF Hessian	122/134 (148)	108/119 (147)	93/107 (143)
UHF	1/15 (20)	1/13 (22)	1/18 (26)
Conv. MP2	16/28 (36)	13/24 (33)	7/21 (31)
Direct MP2	17/53 (60)	14/45 (54)	7/43 (50)
MP2 Gradient	62/90 (112)	55/79 (105)	101/119 (359)
MP2 Hessian	605/658 (693)	585/609 (1127)	1358/1379 (2987)
MP4(SDTQ)	393/405 (542)	372/383 (577)	354/368 (565)
SDCI	33/345 (648)	27/282 (888)	18/174 (771)
CCSD	53/593 (2283)	42/474 (3145)	35/364 (2797)
CCSD(T)	954/969 (2676)	831/842 (4455)	
QCISD	40/410 (858)	33/345 (1227)	
QCISD(T)	871/883 (1210)	681/691 (1681)	
CASSCF	12/249 (930)	8/179 (308)	
CAS-CI	NA	NA	NA
SVWN (LSD)	NA	3/20 (27)	
BLYP (NLSLSD)	NA	4/18 (24)	

Method	MOLPRO (94.8)	GAMESS-US 11/17/94	HONDO (8.3)
Conv. RHF	2/16 (19)	1/8 (11)	
Direct RHF	NA	5/42 (43)	
RHF Gradient	16/32 (35)	8/16 (20)	
RHF Hessian	NA	162/170 (179)	
UHF	1/17 (19)	1/11 (15)	
Conv. MP2	<1/16 (19)	4/12 (16)	
Direct MP2	NA	NA	NA
MP2 Gradient	NA	NA	NA
MP2 Hessian	NA	NA	
MP4(SDTQ)	11/27 (30)	NA	
SDCI	1/23 (26)	16/155 (198)	
CCSD	1/28 (31)	NA	NA
CCSD(T)	20/36 (40)	NA	
QCISD	1/25 (28)	NA	NA
QCISD(T)	19/35 (38)	NA	
CASSCF	1/20 (22)	10/110 (135)	
CAS-CI		NA	
SVWN (LSD)	2/20 (23)	NA	
BLYP (NLSLSD)	2/22 (25)	NA	

Table 15. IBM RS/6000 590 Timings (cont.)

<u>Method</u>	<u>GAMESS-UK</u>	<u>SUPERMOLECULE</u>	<u>ACES II</u>
Conv. RHF		1/18 (20)	
Direct RHF		6/87 (89)	NA
RHF Gradient		64/82 (84)	
RHF Hessian		NA	
UHF		NA	
Conv. MP2		NA	
Direct MP2		74/161 (163)	NA
MP2 Gradient		NA	
MP4(SDTQ)		NA	
SDCI		NA	
CCSD		NA	
QCISD		NA	
CASSCF		NA	NA

Table 15. IBM RS/6000 590 Timings (cont.)

Ethylene, 16 electrons, 1A_g (D_{2h}), Basis Set=cc-pVTZ
 (116 basis functions, 5-term d's, 7-term f's)

Method	Gaussian 92 (E)	Gaussian 92/DFT
Conv. RHF	8/82 (90)	7/69 (76)
Direct RHF	24/236 (244)	19/192 (206)
RHF Gradient	76/158 (167)	63/132 (140)
RHF Hessian	905/987 (1013)	1052/1121 (1166)
UHF		7/85 (93)
Conv. MP2		193/262 (289)
Direct MP2		176/368 (383)
MP2 Gradient		443/705 (747)
MP4(SDTQ)		
SDCI		
CCSD		
CCSD(T)		
QCISD		
QCISD(T)		
CASSCF		
SVWN (LSD)	NA	
BLYP (NLSD)	NA	

Method	MOLPRO (94.8)	GAMESS-US 7/17/93	HONDO (8.3)
Conv. RHF	10/102 (110)		
Direct RHF	NA		
RHF Gradient	NA		
RHF Hessian	NA		
UHF	10/104 (121)		
Conv. MP2	1/103 (110)		
Direct MP2	NA	NA	NA
MP2 Gradient	NA	NA	NA
MP4(SDTQ)	44/146 (154)	NA	
SDCI	3/120 (134)		
CCSD		NA	NA
CCSD(T)			
QCISD		NA	NA
QCISD(T)			
CASSCF			
SVWN (LSD)			
BLYP (NLSD)			

Table 15. IBM RS/6000 590 Timings (cont.)

Ethylene, 16 electrons, 1A_g (D_{2h}), Basis Set=6-311++G(3df,3pd)
 (150 functions, 5-term d', 7-term f's)

Method	Gaussian 92 (E)	Gaussian 92/DFT
Conv. RHF	12/116 (152)	
Direct RHF		
RHF Gradient		
RHF Hessian		
UHF		
Conv. MP2		
Direct MP2		
MP2 Gradient		
MP4(SDTQ)		
SDCI		
CCSD		
CCSD(T)		
QCISD		
QCISD(T)		
CASSCF		
SVWN (LSD)	NA	
BLYP (NLSD)	NA	

Method	MOLPRO (92.3)	GAMESS-US 7/17/93	HONDO (8.3)
Conv. RHF			
Direct RHF	NA		
RHF Gradient			
RHF Hessian	NA		
UHF			
Conv. MP2			
Direct MP2	NA	NA	NA
MP2 Gradient	NA	NA	NA
MP4(SDTQ)		NA	
SDCI			
CCSD		NA	NA
CCSD(T)			
QCISD		NA	NA
QCISD(T)			
CASSCF			
SVWN (LSD)			
BLYP (NLSD)			

Table 15. IBM RS/6000 590 Timings (cont.)

18-crown-6, C₁₂H₂₄O₆, 144 electrons, C_i, Basis Set=3-21G
(210 functions)

<u>Method</u>	<u>Gaussian 92 (E)</u>	<u>Gaussian 92/DFT (G)</u>	<u>Gaussian 94 (C)</u>
Direct RHF	42/539 (558)	32/419 (474)	35/450 (484)
RHF Gradient	216/755 (774)	185/604 (648)	176/626 (673)
RHF Hessian	19665/20204 (20557)	14925/15344 (15747)	13287/13737 (13977)
Conv. RHF		65/784 (5166)	
Conv. MP2			
Direct MP2			
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

<u>Method</u>	<u>GAMESS-US 6/17/92</u>	<u>HONDO (8.1)</u>	<u>MOLPRO (92.3)</u>
Direct RHF			
RHF Gradient			
RHF Hessian			NA
Conv. RHF			
Conv. MP2			
Direct MP2			NA
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

Table 15. IBM RS/6000 590 Timings (cont.)

18-crown-6, C₁₂H₂₄O₆, 144 electrons, C_i, Basis Set=6-31G**
(390 functions)

Method	Gaussian 92 (E)	Gaussian 92/DFT (G)	Gaussian 94 (C)
Direct RHF	323/4197 (4415)	260/3383 (3514)	270/3509 (3630)
RHF Gradient		1462/4755 (4899)	1461/4970 (5104)
RHF Hessian		203730/207113 (209438)	123066/126575 (131317)
Conv. RHF			
Conv. MP2			
Direct MP2			
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

Method	GAMESS-US 6/17/92	HONDO (8.1)	GAMESS-UK (2)
Direct RHF			
RHF Gradient			
RHF Hessian			
Conv. RHF			
Conv. MP2			
Direct MP2			
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

Method	DISCO (1.82)	ACES II	MOLPRO (92.3)
Direct UHF		NA	
RHF Gradient			
RHF Hessian	NA		NA
Direct RHF			
Conv. MP2	NA		
Direct MP2		NA	NA
MP2 Gradient	NA		
MP4(SDTQ)	NA		
SDCI	NA		
CCSD	NA		
QCISD	NA		
CASSCF	NA	NA	

Table 15. IBM RS/6000 590 Timings (cont.)

18-crown-6, C₁₂H₂₄O₆, 144 electrons, C_i, Basis Set=aug-cc-pVDZ
(606 functions)

<u>Method</u>	<u>Gaussian 92 (E)</u>	<u>Gaussian 92/DFT</u>	<u>MOLPRO (92.3)</u>
Direct RHF	9210/147360 (147563)		
RHF Gradient			
RHF Hessian			NA
Conv. RHF			
Conv. MP2			
Direct MP2			NA
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

<u>Method</u>	<u>GAMESS-US 6/17/92</u>	<u>HONDO (8.3)</u>	<u>GAMESS-UK (2)</u>
Direct RHF			
RHF Gradient			
RHF Hessian			
Direct MP2			
MP2 Gradient			
MP4(SDTQ)			
SDCI			
CCSD			
QCISD			
CASSCF			

<u>Method</u>	<u>DISCO (1.82)</u>	<u>ACES II</u>
Conv. RHF		
Direct RHF		NA
RHF Gradient		
RHF Hessian	NA	
UHF	NA	
Conv. MP2	NA	
Direct MP2		NA
MP2 Gradient	NA	
MP4(SDTQ)	NA	
SDCI	NA	
CCSD	NA	
QCISD	NA	
CASSCF	NA	NA

Table 15. IBM RS/6000 590 Timings (cont.)

- (a) All times are in seconds. CPU times are the sum of the "user + system" contributions. Wall clock times are given in parentheses. For the iterative methods (RHF, UHF, SD-CI, QCISD and CASSCF) each entry consists of a trio of numbers: "CPU-time-per-iteration/total-CPU (total-wall-clock)". The "CPU-time-per-iteration" for the conventional SCF methods was defined as the total run time (integrals + SCF) divided by the number of iterations. These values are intended to facilitate comparison with direct HF methods. For other methods the leftmost entry corresponds to the incremental time for the method. For example, the MP2 entry preceding the slash is the total run time minus the time needed for the HF step.
- Calculations were performed on a machine with 128 MB of memory and one 2 GB IBM SCSI 2 disk running under AIX 3.2 with Release 2.3.5 of XLF Fortran. Runs were made on an otherwise quiet system.
- NA:** not available with this program.
FTC-ND: Failed to complete - not enough disk space.
FTC-unknown: Failed to complete for unknown reasons.
- SCF calculations were converged to approximately 15 digits after the decimal point (8 digits in the density).
- (b) The ethylene UHF calculation treated the $\pi \emptyset \pi^* ({}^3B_{1u})$ state. The ethylene ground state is 1A_g . MP2, MP4, CISD and QCISD calculations involved all electrons, i.e., there were no "core" electrons. The CAS configuration list contains 8 CSF's in D_{2h} symmetry and was generated with 4 electrons in 4 orbitals ($3_{ag}, 1b_{3u}, 1b_{2g}, 2b_{1u}$). This configuration list is sufficient to allow ethylene to dissociate into two singlet methylenes. The time reported includes the time required to compute the integrals and solve the CAS equations using the canonical RHF orbitals as the starting guess. The default INDO initial guess used by Gaussian for ethylene's open shell calculations did not pick up the $\pi \emptyset \pi^* ({}^3B_{1u})$ state. If the ordering of the initial guess orbitals was corrected using an ALTER command the calculation with Gaussian 90 died with a complaint that symmetry was being broken. Thus, it was necessary to run these calculations with the NOSYMM option, which ignored the available D_{2h} symmetry. Gaussian 92 fixed this problem with the UHF benchmark and it was run in full D_{2h} symmetry.